



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

where g =gravity=32, l =length to center of oscillation= $\frac{2}{3}a$, h =height of zero velocity= $2l$ without a sensible error, t =time, θ =angle counted from the down point. Reducing, we have,

$$t = \frac{1}{2} \sqrt{\left(\frac{l}{g}\right)} \int_{\frac{1}{2}\pi}^{\pi-\phi} \frac{d\theta}{\cos \frac{1}{2}\theta} = \sqrt{\left(\frac{l}{g}\right)} \left[\log \tan\left(\frac{1}{4}\pi + \frac{1}{4}\theta\right) \right]_{\frac{1}{2}\pi}^{\pi-\phi}$$

$$= \sqrt{\left(\frac{l}{g}\right)} \log \frac{\cot \frac{1}{4}\phi}{\tan \frac{3}{8}\pi} = \sqrt{\left(\frac{l}{g}\right)} (8.647491).$$

But $l = \frac{2}{3}$ of 100. $\therefore t = \sqrt{\left(\frac{100}{32}\right)} (8.647491) = 12.481$ seconds.

AVERAGE AND PROBABILITY.

139. Proposed by L. C. WALKER, A.M., Professor of Mathematics, Colorado School of Mines, Golden, Col.

Four points are taken at random on the surface of a given sphere; find the average volume of the tetrahedron formed by the planes passing through the points taken three and three.

No solution has been received.

140. Proposed by L. C. WALKER, A.M., Professor of Mathematics, Colorado School of Mines, Golden, Col.

Obtain the average area of a triangle formed by a tangent to the four-cusped hypocycloid and the co-ordinate axes.

Solution by J. E. SANDERS, Hackney, Ohio.

The curve's equation is $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$ The length of the perpendicular from the origin on the tangent is

$$\frac{xdx - ydy}{\sqrt{(dx^2 + dy^2)}} = \frac{x^{\frac{2}{3}} + y^{\frac{2}{3}}}{\sqrt{(y^{-\frac{2}{3}} + x^{-\frac{2}{3}})}} = \frac{2}{3} \sqrt{(axy)}.$$

The equation to the tangent is $x'/x^{\frac{1}{3}} + y'/y^{\frac{1}{3}} = a^{\frac{1}{3}}$, and its length between the axis= $a^{\frac{2}{3}} \sqrt{(x^{\frac{2}{3}} + y^{\frac{2}{3}})} = a$.

\therefore The area of the triangle is $A = \frac{1}{2} a^{\frac{2}{3}} \sqrt{(axy)} = \frac{1}{2} a^2 \sin \theta \cos \theta$, if $x = a \cos^3 \theta$, and $y = a \sin^3 \theta$.

Then, if the tangent is through any point on the curve, the average area is

$$\Delta = \frac{1}{s} \int A ds.$$

But $ds = a^{\frac{1}{3}} x^{-\frac{1}{3}} dx = 3a \sin \theta \cos \theta d\theta$ and $s = \frac{2}{3} a$.

$\therefore \Delta = a^2 \int_0^{\frac{1}{2}\pi} \sin^2 \theta \cos^2 \theta d\theta = \frac{1}{16} \pi a^2$, or, if the tangent is to be a random line, the average area is